You will be advised to have your baby before 41 weeks of pregnancy and if there are pregnancy complications affecting either you or your baby, your doctor may recommend birth earlier than this.

What happens in labour?

It is important that your blood glucose level is controlled during labour and birth and it should be monitored to ensure it is not too high. You may be advised to have an insulin drip to help control your blood glucose level.

What happens after my baby is born?

- Your baby will stay with you unless they need extra care. You can usually have skin-to-skin contact with your baby straight away if you choose this. Occasionally they may need to be looked after in a neonatal unit if they are unwell or need extra support.
- Your baby should have their blood glucose level tested a few hours after birth to make sure that it is not too low.
- Gestational diabetes usually goes away after birth and therefore you will be advised to stop taking all diabetes medications immediately after your baby is born. Before you go home, your blood glucose level will be tested to make sure that it has returned to normal.
- You should be offered a fasting blood glucose test 6–13 weeks after the birth of your baby. A small number of women continue to have high blood glucose levels and will be offered further tests for diabetes.
- You should be offered information about your lifestyle, including diet, exercise and watching your weight, to reduce your chance of type 2 diabetes in the future.
- Up to 50% of women who have had gestational diabetes develop type 2 diabetes within the following 5 years. You will therefore be advised to have a test for this every year.

What are my options for feeding my baby?

- Breastfeeding is safe if you have gestational diabetes and your healthcare team will support you in feeding your baby.
- Whichever way you choose to feed your baby, you should start feeding as soon as possible after birth, and then every 2–3 hours to help your baby's blood glucose stay at a safe level. Babies born to mothers with gestational diabetes have a high risk of low sugar levels after birth, so you may be advised to hand express and give your baby this early breast milk (also called colostrum) in addition to breastfeeding directly.

What do I need to know about future pregnancies?

Having a healthy weight, eating a balanced diet and taking regular physical exercise before you become pregnant can reduce your risk of developing gestational diabetes again.

As soon as you find out you're pregnant, contact your doctor for advice about your antenatal care as there is a chance you may develop gestational diabetes again (more than 1 in 3 women will get gestational diabetes again).

Emotional support

Having a healthy weight, eating a balanced diet and taking regular physical exercise before you become pregnant can reduce your risk of developing gestational diabetes again.

As soon as you find out you're pregnant, contact your doctor for advice about your antenatal care as there is a chance you may develop gestational diabetes again (more than 1 in 3 women will get gestational diabetes again).

REFERENCES : NICE GUIDELINE 2020

LOCATION

CONSULT OUR SPECIALISTS
+971 26 11 33 33 24x7
Shabia 9, Mussafah, P.O.Box: 91250, Abu Dhabi,UAE
info@millenniumhospital.ae
www.millenniumhospital.ae



PATIENT EDUCATION LEAFLET

Follow us



Gestational Diabetes

مستشفى ميلينيوم
MILLENNIUM HOSPITAL
Specialized in
WOMEN & CHILD CARE
عنوان جديد للعناية الشاملة بالأم والطفل

What is gestational diabetes?

Diabetes that develops during pregnancy is known as gestational diabetes. It happens because your body cannot produce enough insulin (a hormone that helps to control blood glucose) to meet the extra needs of pregnancy.

Gestational diabetes usually starts in the middle or towards the end of pregnancy

How common is gestational diabetes?

Gestational diabetes is common. It affects at least 4–5 in 100 women during pregnancy.
You are more likely to develop gestational diabetes if you have any of the following risk factors:

- your body mass index (BMI) is higher than 30
- you have previously given birth to a baby weighing 4.5 kg (10 lbs) or more
- you have had gestational diabetes before
- you have a parent, brother or sister with diabetes
- your family origin is South Asian, Chinese, African-Caribbean or Middle Eastern.

How will I be checked for gestational diabetes?

If you have any of the above risk factors, you should be offered a glucose test during your pregnancy. This may be a blood test in early pregnancy and/or a glucose tolerance test (GTT) when you are between 24 and 28 weeks pregnant.

A GTT involves fasting overnight (not eating or drinking anything apart from water):

- In the morning, before breakfast, you will have a blood test. You are then given a glucose drink.
- The blood test is repeated 1–2 hours later to see how your body reacts to the glucose drink.

During your routine pregnancy care, your urine is tested for glucose. If glucose is present in your urine, then your healthcare team may recommend that you have a GTT.

What does gestational diabetes mean for me and my baby?

Most women who develop gestational diabetes have healthy pregnancies and healthy babies but occasionally gestational diabetes can cause serious problems, especially if it is not recognised or treated.
If your blood glucose levels are high, the chances of you having an induced labour or a caesarean birth are increased.

The risks to your baby are:

- being bigger than average
- shoulder dystocia (where your baby's shoulder gets stuck during birth)
- stillbirth or the baby dying at or around the time of birth. This is uncommon.
- needing additional care once they have been born, possibly in a neonatal unit
- being at greater risk of developing obesity and developing type 2 diabetes in later life.

Controlling your levels of blood glucose during pregnancy and labour reduces the chances of these complications for you and your baby.

What extra care will I need during pregnancy?

If you are diagnosed with gestational diabetes, you will be under the care of a specialist team and will be advised to have your baby in a hospital with facility for neonatal care.
Your team may include a doctor specialising in diabetes, an obstetricianand dietitian. You should start receiving extra antenatal care as soon as your gestational diabetes is diagnosed.

Healthy eating and exercise

The most important treatment for gestational diabetes is a healthy eating plan and exercising regularly. Walking for 30 minutes after a meal can help with controlling your blood glucose levels. Gestational diabetes usually improves with these changes.

Monitoring your blood glucose

After you have been diagnosed with gestational diabetes, you will be shown how to check your blood glucose levels and told what your ideal level should be. If it does not reach this level with healthy eating and exercise, or if an ultrasound scan shows that your baby is larger than expected, you may need to take tablets or give yourself insulin injections. If your glucose level is very high at the time of diagnosis, then you may be offered treatment straight away, in addition to making changes to your diet and exercise.

Monitoring your baby

You should be offered extra ultrasound scans to monitor your baby's growth more closely
Advice and information
During your pregnancy, your doctor will give you information and advice about:

- planning birth, including timing and types of birth, pain relief and changes to your medica tions during labour and after your baby is born
- looking after your baby following birth
- care for you after your baby is born including contraception.

Will I need treatment?

Some women with gestational diabetes will need to take tablets and/or have insulin injections to control their blood glucose during pregnancy. Your specialist team will advise you what treatment is best for you and your baby.
If you do need insulin, your specialist healthcare team will explain exactly what you need to do. This will include showing you how to inject yourself with insulin, how often to do it and when you should check your blood glucose levels.

What are my birth options?

You will have discussions about your options for birth with your doctor throughout your pregnancy. Your options include waiting for labour to start, having an induction of labour or having a planned caesarean birth. This will depend on your individual circumstances and preferences and your doctor will discuss the risks and benefits of each option with you.